## Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

## **Listing of Claims:**

1. (withdrawn) A method for preparing monoglyceride sulfonate, which comprises:

a step of neutralizing a fatty acid derived from an animal oil such as tallow and lard, or from a plant oil selected from a group consisting of coconut oil, lauric acid, palm oil, and palm kernel oil, with an alkali metal solution in a solvent to prepare a single or mixed alkali metal salt of a fatty acid; and

a step of reacting the salt with a compound represented by the following Chemical Formula 2:

## Chemical Formula 1

Chemical Formula 2

CICH2CH(OH)CH2SO3-M

wherein R is a  $C_7$  to  $C_{19}$  saturated or unsaturated aliphatic hydrocarbon radical, and M is sodium or potassium.

2. (withdrawn) The method for preparing monoglyceride sulfonate according to claim 1, wherein the compound represented by Chemical Formula 2 is prepared by reacting epichlorohydrin with sodium sulfite, sodium bisulfite, or sodium metabisulfite.

- 3. (withdrawn) The method for preparing monoglyceride sulfonate according to claim 1, wherein the alkali metal salt of a fatty acid and the compound represented by Chemical Formula 2 are reacted in the reaction equivalent ratio of 1:0.05 to 1.2.
- 4. (withdrawn) The method for preparing monoglyceride sulfonate according to claim 1, wherein the solvent is water, or a mixture of water and a low alcohol.
- 5. (withdrawn) A cleansing agent for a human body prepared by the method according to claim 1.
- 6. (withdrawn) A soft soap composition comprising:

50 to 90 parts by weight of a mixture of monoglyceride sulfonate represented by the following Chemical Formula 1, which contains more than 60 wt % of lauric acid and myristic acid, and a fatty acid soap;

1 to 12 parts by weight of a fatty acid; and

1 to 25 parts by weight of a binder (plasticizer) or an excipient:

## Chemical Formula 1

wherein R is a  $C_7$  to  $C_2$ , alkyl; and M is sodium, potassium, triethanolamine, or ammonia.

- 7. (withdrawn) The soft soap composition according to claim 6, wherein the mixing ratio of the monoglyceride sulfonate and the fatty acid soap is from 1:0.3 to 0.03:1.
- 8. (withdrawn) The soft soap composition according to claim 6, wherein the content of the lauric acid and the myristic acid is over 70 wt %.

- 9. (withdrawn) The soft soap composition according to claim 6, which further comprises 1 to 25 parts by weight of a surfactant.
- 10. (currently amended) A method for preparing a soft soap containing salt ( $\underline{NaCl}$ ), which comprises:
- (a) a step of neutralizing a  $C_8$  to  $C_{22}$  saturated or unsaturated fatty acid with caustic soda to obtain a fatty acid sodium salt represented by the following Chemical Formula 3a; and
- (b) a step of reacting the fatty acid sodium salt with 3-chloro-2-hydroxypropanesulfonic acid sodium salt (SCHS) represented by the following Chemical Formula 2a in a solvent, wherein the soft soap contains 2 to 15 wt % of said salt (NaCl) the salt:

Chemical Formula 3a

RCOO Na

wherein R is a  $C_7$  to  $C_{21}$  saturated or unsaturated aliphatic hydrocarbon;

Chemical Formula 2a

ClCH<sub>2</sub>CH (OH) CH<sub>2</sub>SO<sub>3</sub>Na

wherein said 2 to 15 wt% of salt (NaCl) is formed as a product of the reaction of said fatty acid sodium salt with said 3-chloro-2-hydroxypropanesulfonic acid sodium salt (SCHS);

and wherein the fatty acid salt contains more than 60 wt% of lauric acid and myristic acid.

- 11. (currently amended) The method for preparing a soft soap containing salt (NaCl) according to claim 10, wherein in the step (a) the  $C_8$  to  $C_{22}$  saturated or unsaturated fatty acid is used alone or in combination.
- 12. (currently amended) The method for preparing a soft soap containing salt (NaCl) according to claim 10, wherein in the step (b) reaction equivalent ratio of the fatty acid sodium salt to the 3-chloro-2-hydroxypropanesulfonic acid sodium salt is from 1:0.1 to 1:1.2.
- 13. (canceled)